Gravitation

We have already talked about the force of gravity:



The main assumption needed to use the above expression for the force of gravity is that the <u>person needed to be very close to the surface of the earth</u>.

However, the force of gravity acts beyond just the surface of the earth. The force of gravity is responsible for:

- Keeping the moon in orbit around the earth
- Keeping the earth in orbit around the sun
- Keeping the solar system as part of the galaxy
- etc

Gravity acts between all things with mass.

Newton's Law of Universal Gravitation

The gravitational force between any two bodies of mass m_1 and m_2 , separated by a distance r, is given by:



$$F_{G} = \frac{Gm_{1}m_{2}}{r^{2}}$$

$$G = 6.67 \times 10^{-11} \text{ N} \cdot \text{m}^2 / \text{kg}^2$$

Note:

- 1) Object m_1 attracts object m_2 with force F_{12}
- 2) This force is attractive, and is directed towards m_1
- 3) Object m_2 attracts object m_1 with force F_{21}

 $\left| \mathbf{F}_{12} \right| = \left| \mathbf{F}_{21} \right| = \mathbf{F}_{G}$

4) This force is also attractive and directed towards m_2

Gravity Near the Surface of the Earth



 F_{EP} = Force of earth on the person

 F_{PE} = Force of person on the earth

$$\left| \mathbf{F}_{\mathrm{EP}} \right| = \left| \mathbf{F}_{\mathrm{PE}} \right| = \mathbf{F}_{\mathrm{G}}$$

If the person always stays very close to the surface of the earth, then r_E can be approximated as a constant.

The force of gravity (due to the earth) on an object with mass m very near the earth's surface:

$$F_{\rm G} = \frac{{\rm Gm}_{\rm E}m}{{\rm r}_{\rm E}^{2}} = \left(\frac{{\rm Gm}_{\rm E}}{{\rm r}_{\rm E}^{2}}\right)m = mg$$

$$\left(\frac{\mathrm{Gm}_{\mathrm{E}}}{\mathrm{r}_{\mathrm{E}}^{2}}\right) \equiv \mathrm{a}_{\mathrm{g}} = 9.83 \mathrm{m/s}^{2}$$

Lecrture 10: Gravitation

Interesting things about Gravity



- G is a measure of the "strength" of gravity. It is very very small.
- So the force of gravity will be large only when at least one of the two masses in the above expression are large.
- But the force of gravity has <u>unlimited range</u>. F becomes zero only in the limit that r goes to infinity.
- This last statement implies that <u>you are currently exerting a (small)</u> <u>force on galaxies far, far away</u>.
- The above expression is only strictly true for "pointlike" particles. How is it that we can use it for "extended" objects like the earth, a car, ourselves...?

Principle of Superposition

Principle of superposition: The net effect is the sum of individual effects.



The total gravitational force on particle m_1 due to particles m_2 , m_3 , and m_4 , is the vector sum of the gravitational forces:

$$\vec{F}_1 = \vec{F}_{21} + \vec{F}_{31} + \vec{F}_{41}$$

Or more generally for "n" particles:

$$\vec{F}_1 = \sum_{i=2}^n \vec{F}_{i1}$$

If we want to know the force that a real object (made up of a continuous distribution of particles) exerts on another particle of mass m, we need to replace the above discrete sum by a continuous integral:

$$F_1 = \int dF = \int \frac{Gm_1}{r^2} dm$$

Example: Force on a Particle due to a Ring:

A particle of mass m_1 is located a distance z away from a thin ring of mass m_2 , along its axis. The radius of the ring is R.

Determine the magnitude and direction of the gravitational force of the ring, on the particle m_1 .



Example: Force on a Particle due to a Ring:



First consider a small element dm_2 of the ring.

 dF_{Gz} : This is the z-component of the force of gravity due to mass element dm_2 .

Notice that the component of the force of gravity *perpendicular* to this will be canceled by a mass element on the opposite side of the ring.

The z-component of the force of gravity exerted on mass m_1 by mass element dm_2 is:

$$dF_{Gz} = \left(\frac{Gm_1 dm_2}{r^2}\right) \cos \hat{e}$$

Lecrture 10: Gravitation

Example: Force on a Particle due to a Ring:

First consider a small element dm_2 of the ring.



The force of gravity exerted on mass m_1 by the whole ring is then:

$$F_{Gz} = \int dF_{Gz} = \frac{z Gm_1 m_2}{(z^2 + R^2)^{3/2}}$$

Lecrture 10: Gravitation

Ex: Force on mass due to a thin Ring

NOTE:

 If the mass m₁ is located at z=0, the gravitational force goes to zero.
Newton showed using similar methods that a uniform spherical shell of mass m attracts particles outside of the shell as if all of the shell's mass were located at its center. The same can be demonstrated for a uniform sphere.
If a particle is located anywhere inside a uniform shell, the net gravitational attraction on it is zero.

Example: Gravitational Attraction between Two Spheres



Both spheres have mass m=100kg, and are separated by a distance of 1.00m.

$$F_{\rm G} = \frac{{\rm Gm}_1 {\rm m}_2}{r^2} = \frac{(6.67 \times 10^{-11})(100)(100)}{(1.00)^2}$$
$$F_{\rm G} = 6.67 \times 10^{-7} \,\rm N$$

This is small. About the weight of a baby flea. Will not pull a lineman offsides.

Example: Gravitational Attraction between The Earth and the Moon



This is big. Equivalent to the thrust of 6 million Saturn 5 moon rockets.