

## Physics 880.06: Problem Set 6

Due Friday, November 8, 2002

1. Ashcroft and Mermin, Chapter 9, Problem 3.
2. Consider the tight-binding approximation for an  $s$ -band in a triangular lattice with lattice constant  $a$ . Let the normalized atomic wave function be denoted  $\phi(\mathbf{r})$ . Assume that the hopping matrix element  $H_{\mathbf{R},\mathbf{R}'} \equiv \int_d^3 r \phi(\mathbf{r} - \mathbf{R}) H \phi(\mathbf{r} - \mathbf{R}') = -t$  ( $t > 0$ ) if  $\mathbf{R}$  and  $\mathbf{R}'$  are nearest neighbor Bravais lattice vectors,  $H_{\mathbf{R},\mathbf{R}'} = \epsilon_0$  if  $\mathbf{R} = \mathbf{R}'$ , and  $H_{\mathbf{R},\mathbf{R}'} = 0$  otherwise.
  - (a) Find the band energies  $E(\mathbf{k})$  in terms of  $t$ ,  $a$ , and  $\epsilon_0$ .
  - (b). Show that near the bottom of the band  $E(\mathbf{k}) = E_{min} + \hbar^2 k^2 / (2m^*)$ , and find  $E_{min}$  and  $m^*$  in terms of  $\epsilon_0$ ,  $t$ , and  $a$ .
  - (c). What is the band width, in terms of  $t$ ,  $\epsilon_0$ , and  $a$ ?
3. *For edification only; not to be turned in.* Consider a layered dielectric consisting of alternate layers of thickness  $d_1$  and  $d_2$ , having dielectric constants  $\epsilon_1$  and  $\epsilon_2$ , and consider a linearly polarized plane electromagnetic wave propagating perpendicular to the layers. Let this be denoted the  $z$  direction.
  - (a). Write down the form of the electric field, given that it must satisfy Bloch's theorem. Write down the corresponding magnetic field  $\mathbf{B}$ . (Assume that the relative permeability  $\mu = 1$  in both media.)
  - (b). What are the boundary conditions on  $\mathbf{E}$  and  $\mathbf{B}$  at  $z = d_1$  and  $z = d_1 + d_2$ ?
  - (c). Using (a) and (b), find a determinantal equation which gives a relationship between the Bloch vector  $k$  and the frequency  $\omega$ . Don't try to solve this equation.